

72215
Aphanitic Impact Melt Breccia
379.2 grams




Figure 1: Boulder #1 at station #2 on the slope of South Massif, Apollo 17. AS17-137-20900. Boulder is ~2 meters across. Lines roughly indicate apparent layering in boulder (Wolfe et al. 1981).

Introduction

Lunar sample 72215 was collected from the side of a layered boulder #1 at station 2 located on the bottom slope of the South Massif and within the landslide material at Apollo 17 Taurus-Littrow (Schmitt 1973, Wolfe et al. 1981). Samples 72235, 72255 and 72275 are from other layers in this boulder and soils 72220, 72240 and 72260 are from the fillet surrounding the boulder. The boulder had a prominent layering with multi-layered clasts weathering out as “knobs” (figure 1). 72215 was sampled as one of the largest of these “knobs” (~10 cm).

The “knobs” are intricate swirls of light and dark, interlayered, crushed highland materials produced in some, as yet, unexplained fluidized impact mechanism. The dark portions are finer-grained and the lighter are coarser-grained – but both portions are found to have similar bulk composition. Two of these “knobs” were given special names: Marble Cake (72275) and Dying Dog (72235) by the Consortium Indomitable who first studied them. Aspects of 72215 are similar.

The sample is a coherent, clast-rich, impact melt breccia. It is aluminous and feldspathic and contains




Figure 2: Exterior surface of 72215 showing patina and zap pits. Cube is 1 cm. NASA S73-23569.

numerous small patches of “granitic” material. It has been preliminary dated at 3.83 b.y. and has an exposure age of 42 m.y.

Petrography

Ryder (1993) provides a complete review of the work accomplished on 72215. Briefly, 72215 has a prominent layering that has been described by Ryder et al. (1974) and Stoeser et al. (1975) as separate “domains” – seen in figure 3 and 4.

Domains 1, 2 and 3 are similar in chemical composition, but differ in color and average





Figure 4: Photographs of thin sections 72215, 180, 183 and ,190, along with sketch maps (continued on next page). Together these sections provide an almost complete microscopic view across the whole sample (about 6 cm). The “knob” is the portion to the left (see figure 14). The different “domains” are given numbers (circled). From Stoeser et al. 1974.




Figure 3: Freshly broken surface of 72215, illustrating banding and clastic nature. Cube is 1 cm. NASA S73-23569. GA is the position of the "knob".

grain size. They are composed of angular and rounded mineral grains and small clasts with a seriate grain size down to about 20 microns – in areas, the matrix is less than 1 micron. Analyses show it is feldspathic.

Domain 4 consists of crushed, fine-grained feldspathic granulite strung out in lenses and mixed with matrix for other domains. Following cataclasis, annealing was sufficient to eliminate porosity.




Figure 4 continued




Figure 5: Composition of pyroxene in matrix 72215 (Ryder et al. 1975).




Figure 6: Chemical composition of pyroxene in granite clasts in 72215 (Ryder et al. 1975).




Figure 7: K, Na, Ca diagram for feldspar grains found in "granitic" patches in 72215, 72235 and 72255 (Ryder et al. 1975).




Figure 8: Poikilitic ANT clast in 72215, 108. Plagioclase and olivine chadocysts enclosed in a large orthopyroxene oikocryst. Field of view is about 700 microns. From Stoeser et al. 1974.




Figure 9: Evidence of cohenite at boundary of Ni, Co, Fe grain in 72215 (Goldstein et al. 1976).

Composition of ternary feldspar in 72215.

(from Ryder et al. 1975)

SiO ₂	53.28	53.14	53.04	51.09	52.85	54.37
Al ₂ O ₃	30.1	30.02	28.77	29.97	28.05	28.89
FeO	0.2	0.22	0.23	0.21		
MgO	0.04	0.02	0.05	0.04		
CaO	13.82	13.78	12.97	13.98	13.36	13.51
Na ₂ O	2.91	2.66	2.67	2.38	2.17	1.27
K ₂ O	1.21	1.53	1.57	2.09	2.43	4.44
BaO	0.13	0.12	0.07	0.13		0.08
total	101.7	101.5	99.39	99.88	98.87	102.57
electron microprobe						

Domain 5 is darker than the others and more vesicular. It has an abundance of granitic patches or clasts (up to 23%!). Broad beam analyses show it has higher K₂O.

Domain 6 (GA) is white, coarse-grained and cataclasized. It is made up of relic areas of poikilitic granulite and crushed equivalent. Equidimentional chadocysts of plagioclase (An₉₀₋₉₆) are embedded in large pyroxene oikocrysts (En₇₅Wo₄).

Significant Clasts

Granite

The “granitic” clasts in boulder 1, including those in 72215, were analyzed and described by Ryder et al. (1975) and Stoeser et al. (1995). They have high silica (>75%) and high K₂O contents (~6%)(table 1). They are made of various intergrowths of silica, Ba,K feldspar, ternary feldspar (figure 7) and Fe-rich pyroxene (figure 6). Compston et al. (1975) determined a “mixing line” with apparent age of ~ 4 b.y. for these Rb-rich “granitic” assemblages (figure 11).

GA ,76

The white clast marked as GA in the figures 3, 4 and 14 is a cataclastic, poikilitic anorthositic norite. The analysis show that it is low in trace elements and high in alumina (Blanchard et al. 1974). It is also known as “domain 6”.

Pink Spinel, Troctolite basalt

Small clasts of basalt containing Mg-spinel and/or olivine have been reported (Table 3).




Figure 10: Normalized rare-earth-element diagram for 72215 (data from Blanchard et al. 1975 and Palme et al. 1978).

Chemistry

Palme et al. (1978), Blanchard et al. (1975), Morgan et al. (1975) have analyzed 72215 (table 1, figure 10). In addition, Stoeser used the broad beam electron probe to obtain the composition of the various domains (table 3). Everything is high alumina and there is not much different about the various domains – other than grain size and appearance.

Morgan et al. (1975) found that meteoritic siderophiles (Ir, Re, Os etc.) have different ratios (figure 11) from other lunar breccias and discuss this in context to the Serenitatis ejecta.

Radiogenic age dating

Schaeffer et al. (1982) dated the matrix and various clasts using the Ar/Ar laser technique.

Compston et al. (1975) reported data for an anorthosite clast, the matrix and various granitic clasts (figure 12).

Mineralogical Mode for 72215

From Stoeser et al. 1974

	Domains 1 + 2	Domain 3	Domain 5
ANT breccias	8.2 %	6.7	7.8
Granulitic ANT	19.8	23.5	25
Poikilitic ANT	2.2	6.3	2.6
Coarse ANT	4.5	5.3	3.4
Ultramafic	2.4	1.4	1.7
Basalts	3.6	1.4	0.9
Granite	4.2	11	23.2
Maskelynite	14	10	2.6
Glass	0.4		
Plagioclase	24.3	17.7	21.6
Olivine	5.1	6.7	1.7
Pyroxene	10.6	9.1	9.5
Chromite	0.2		
Silica		0.5	




Figure 11: Chopped 3-element diagram for Ir, Re and Au used by Morgan et al. (1975) to show that station 2 boulder samples at Apollo 17 were different in trace siderophile element content from other Apollo 17 samples.

Nunes and Tatsumoto (1975) studied the U-Th-Pb systematics, but could not obtain an age.

Summary of Age Data for 72215

	Ar/Ar	Rb/Sr
Schaeffer et al.	3.83 – 4.02 b.y.	
Compston et al.		~ 4 b.y.

Cosmogenic isotopes and exposure ages

The cosmic-ray exposure age of 72215 was determined as 41.4 m.y. by ^{81}Kr and 41 m.y. by ^{38}Ar methods (Leich et al. 1975). This is less than the exposure age of samples from the top of the boulder (e.g. 72275), probably because of shielding or erosion effects. Arvidson et al. (1976) have proposed that this is the age of Tyco!

Other Studies

Rare gas contents	Leich et al. 1975
Reflectance spectra figure 13	Adams and Charette 1975
Pb isotopes	Nunes and Tatsumoto 1975
Magnetics	Banerjee et al. 1975
Cohenite	Goldstein et al. 1976
figure 9	




Figure 12: Rb-Sr diagram for "granite" clasts in boulder samples 72215, 72255 and 72275 (Compston et al. 1975).




Figure 13: Reflectance spectra of chips of different lithologies from 72215 (Adams and Charette 1975).

Table 1. Chemical composition of 72215.

reference	Blanchard75	Higuchi75				Palme78	Ryder75					glass
weight	matrix	clast,76	Morgan75	grey matrix	clast4	granite clasts						
SiO ₂ %	45.2	44.7	(b)			46.68	(d)	79.13	78.05	79.58	71.14	74.25
TiO ₂	0.9	0.5	(b)			0.7	(d)	0.1	0.14	0.14	0.31	0.31
Al ₂ O ₃	21.1	27.3	(b)			20.33	(d)	9.44	10.59	10.89	12.48	11.81
FeO	8.43	4.8	(a)			7.81	(d)	0.63	0.17	0.17	0.68	2.01
MnO	0.128	0.067	(b)			0.12	(d)	0.02		0.01	0.02	0.01
MgO	10.7	7.19	(b)			10.33	(d)	0.07			0.12	0.12
CaO	12.1	14.9	(b)			12.5	(d)	1.27	1.39	2.25	3.22	0.53
Na ₂ O	0.52	0.48	(a)			0.47	(d)	0.19	0.16	0.26	0.42	0.03
K ₂ O	0.25	0.11	(b)			0.21	(d)	6	6.66	5.75	6.34	9.55
P ₂ O ₅						0.24	(d)	0.05	0.02	0.02	0.64	0.1
S %						0.032	(d)					0.02
<i>sum</i>												
Sc ppm	18.6	7.68	(a)					18.5	(d)			
V												
Cr								1665	(d)			
Co	27	11.9	(a)					26.9	(d)			
Ni	160	50	(a)	154	146	158	136	56	(c)	170	(d)	
Cu												
Zn				1.8	1.7	1.6	1.8	1.3	(c)			
Ga								3.9	(d)			
Ge ppb				141	117	144	124	42	(c)			
As												
Se				68	49	72	71	15	(c)			
Rb				4.88	2.83	6.65	3.88	1.48	(c)			
Sr								160	(d)			
Y								90	(d)			
Zr								360	(d)			
Nb								25	(d)			
Mo												
Ru												
Rh												
Pd ppb												
Ag ppb				0.46	0.56	0.46	0.47	0.5	(c)			
Cd ppb				3.6	3.8	3	5.3	6.7	(c)			
In ppb												
Sn ppb												
Sb ppb				0.66	0.64	0.92	0.71	0.31	(c)			
Te ppb				3.3	2.4	2.5	4.9	<5.4	(c)			
Cs ppm				0.16	0.12	0.216	0.2	0.05	(c)	0.18	(d)	
Ba								299	(d)			
La	30	7.3	(a)					28.1	(d)			
Ce	77	18.3	(a)					72	(d)			
Pr												
Nd								45	(d)			
Sm	14.5	3.36	(a)					11.2	(d)			
Eu	1.38	1	(a)					1.34	(d)			
Gd								15.5	(d)			
Tb	2.6	0.75	(a)					2.47	(d)			
Dy								17.2	(d)			
Ho								3.77	(d)			
Er												
Tm								1.56	(d)			
Yb	10.3	3.1	(a)					9.29	(d)			
Lu	1.37	0.44	(a)					1.26	(d)			
Hf	9.7	2.4	(a)					9.11	(d)			
Ta	1.4	0.3	(a)					1.14	(d)			
W ppb												
Re ppb				0.37	0.4	0.38	0.28	0.19	(c)			
Os ppb												
Ir ppb				5.2	5.4	5	4	3	(c)	6.1	(d)	
Pt ppb												
Au ppb				2.2	2.1	2.3	1.8	0.8	(c)	4	(d)	
Th ppm	5	1.3	(a)							3.94	(d)	
U ppm				1.36	1.32	1.52	1.62	0.52	(c)	1.16	(d)	

technique: (a) INAA, (b) AA, (c) RNAA, (d) INAA, RNAA, XRF, (e) broad-beam e-probe

Table 2: 72215

Nunes et al. 1975

	U ppm	Th ppm	K ppm	Rb ppm	Sr ppm
	1.408	4.633			
	1.369	5.084			
	1.29	4.635			
	1.179	4.327			
	1.185	4.298			
	1.023	3.768			
Compston et al. 1975				5.02 0.86 4.05 44.19 43.98	139.1 164.6 149.7 162.2 161


Table 3: Broad beam electron probe analyses of domains in 72215.

by Stoeser et al. 1974

	domain 1	domain 2	domain 3	domain 4	domain 5	domain 6	PST	basalt	olivine	basalt
SiO ₂	46.2	45.05	46.75	47.48	48.02	45.81	48.17		43.67	
TiO ₂	0.56	0.6	0.68	0.79	0.66	0.61	0.36		0.2	
Al ₂ O ₃	18.3	19.81	19.64	23.14	18.79	23.93	22.3		18.86	
FeO	8.39	7.32	8.2	6.63	8	5.83	4.73		6.73	
MnO	0.12	0.12	0.11	0.05	0.12	0.05	0.07		0.07	
MgO	11.04	9.71	10.54	6.37	7.32	7.09	9.07		17.24	
CaO	12.01	12.62	12.73	14.76	11.81	14.72	14.44		11.31	
Na ₂ O	0.51	0.56	0.49	0.58	0.77	0.46	0.51		0.39	
K ₂ O	0.21	0.18	0.22	0.34	1.02	0.28	0.35		0.09	

Processing

The original processing of 72215 was documented by Marvin (1974). A sawcut lengthwise through the sample produced a slab (.15) with one side sawn and the other original surface (figure 13). At a later time additional material was provided to Ryder. There are 30 thin sections. The main mass is still intact (as of 2007).






Figure 14: Sawn surface of 72215. Note white cataclastic gabbroic anorthosite (GA) with black rind (the knob). Cube is 1 cm; scale is in mm.. NASA S74-20753. Saw marks obscure texture. Approximate position of set of continuous thin sections (figure 4) is shown.





Figure 15: Map of slab of 72215 - reverse side of figure 14. Position of thin section set (figure 4) is also in reverse order. From Marvin (1974).

References for 72215

- Adams J.B. and Charette M.P. (1975) Spectral reflectance of highland rock types at Apollo 17: Evidence from Boulder 1, Station 2. *The Moon* **14**, 483-489.
- Arvidson R., Drozd R., Guiness E., Hohenberg C., Morgan C., Morrison R. and Oberbeck V. (1976) Cosmic ray exposure ages of Apollo 17 samples and the age of Tyco. *Proc. 7th Lunar Sci. Conf.* 2817-2832.
- Banerjee S.K. and Swits G. (1975) Natural remanent magnetization studies of a layered breccia boulder from the lunar highland region. *The Moon* **14**, 473-481.
- Banerjee S.K., Hoffman K. and Swits G. (1974a) Remanent magnetization directions in a layered boulder from the South Massif. *Proc. 5th Lunar Sci. Conf.* 2873-2881.
- Banerjee S.K., Hoffman K. and Swits G. (1974b) Reversed polarity remanent magnetization in a layered boulder near South Massif (abs). *LS V*, 32-34.
- Blanchard D.P., Haskin L.A., Jacobs J.W., Brannon J.C. and Korotev R.L. (1975) Major and trace element chemistry of Boulder 1 at Station 2, Apollo 17. *The Moon* **14**, 359-371.
- Compston W., Foster J.J. and Gray C.M. (1975) Rb-Sr ages of clasts from within Boulder 1, Station 2, Apollo 17. *The Moon* **14**, 445-462.
- Dalrymple G.B. and Ryder G. (1996) Argon-40/argon-39 age spectra of Apollo 17 highlands breccia samples by laser step heating and the age of the Serenitatis basin. *J. Geophys. Res.* **101**, 26069-26084.
- Goldstein J.I., Hewins R.H. and Romig A.D. (1976a) Carbides in lunar soils and rocks. *Proc. Lunar Sci. Conf. 7th*, 807-818.
- Goldstein J.I., Hewins R.H. and Romig A.D. (1976b) Carbides in lunar soils and rocks (abs). *LS VII*, 310-312.
- Goswami J.N. and Hutcheon I.D. (1975) Cosmic ray exposure history and compaction age of Boulder 1 from Station 2. *The Moon* **14**, 395-405.
- Goswami J.N., Braddy D. and Price P.B. (1976a) Microstratigraphy of the lunar regolith and compaction ages of lunar breccias. *Proc. 7th Lunar Sci. Conf.* 55-74.
- Goswami J.N., Braddy D. and Price P.B. (1976b) Microstratigraphy of the lunar regolith and compaction ages of lunar breccias (abs). *LS VII*, 328-330.
- Higuchi H. and Morgan J.W. (1975) Ancient meteoritic component in Apollo 17 boulders. *Proc. 6th Lunar Sci. Conf.* 1625-1651.
- Hutcheon I.D., Macdougall D. and Stevenson J. (1974b) Apollo 17 particle track studies: surface residence times and fission track ages for orange glass and large boulders. *Proc. 5th Lunar Sci. Conf.* 2597-2608.
- James O.B. (1982) Subdivision of the Mg-suite plutonic rocks into Mg-norites and Mg-gabbronorites (abs). *LPS XIII*, 360-362.
- James O.B. and Flohr M.K. (1983) Subdivision of the Mg-suite noritic rocks into Mg-gabbronorites and Mg-norites. *Proc. 13th Lunar Planet. Sci. Conf., J. Geophys. Res.*, A603-A614.
- Jovanovic S. and Reed G.W. (1975a) Cl and P_2O_5 systematics: Clues to early lunar magmas. *Proc. 6th Lunar Sci. Conf.* 1737-1751.
- Jovanovic S. and Reed G.W. (1975b) Soil breccia relationships and vapor deposits on the moon. *Proc. 6th Lunar Sci. Conf.* 1753-1759.
- Jovanovic S. and Reed G.W. (1975c) History of Boulder 1 at Station 2, Apollo 17 based on trace element interrelationships. *The Moon* **14**, 385-393.
- Jovanovic S. and Reed G.W. (1975d) Studies on regolith processes: Apollo 15 and 17 labile trace element implications (abs). *LS VI*, 451-453.
- Jovanovic S. and Reed G.W. (1980a) Candidate samples for the earliest lunar crust. *Proc. Conf. Lunar Highlands Crust, Geochim. Cosmochim. Acta, Suppl. 12*. Pergamon Press. 101-111.
- Knoll H.-D. and Stöffler D. (1979) Characterization of the basic types of lunar highland breccias by quantitative textural analysis (abs). *LPS X*, 673-675.
- Leich D.A., Kahl S.B., Kirschbaum A.R., Niemeyer S. and Phinney D. (1975a) Rare gas constraints on the history of Boulder 1, Station 2, Apollo 17. *The Moon* **14**, 407-444.
- Macdougall D., Rajan R.S., Hutcheon I.D. and Price P.B. (1973) Irradiation history and accretionary processes in lunar and meteoritic breccias. *Proc. 4th Lunar Sci. Conf.* 2319-2336.
- Macdougall D., Hutcheon I.D. and Price P.B. (1974) Irradiation records in orange glass and two boulders from Apollo 17 (abs). *LS V*, 483-485.

- Marvin U.B. (1974) Appendix B: Specimen 72215 documentation. Consortium Indomitable.
- Marvin U.B. (1975) The Boulder. *The Moon* 14, 315-326.
- Morgan J.W., Ganapathy R., Higuchi H., Krahenbuhl U. and Anders E. (1974a) Lunar basins: Tentative characterization of projectiles, from meteoritic elements in Apollo 17 boulders. *Proc. 5th Lunar Sci. Conf.* 1703-1736.
- Morgan J.W., Ganapathy R., Higuchi H., Krahenbuhl U. and Anders E. (1974b) Lunar basins: Tentative characterization of projectiles, from meteoritic elements in Apollo 17 boulders (abs). *LS V*, 526-528.
- Morgan J.W., Higuchi H. and Anders E. (1975b) Meteoritic material in a boulder from the Apollo 17 site: Implications for its origin. *The Moon* 14, 373-383.
- Nunes P.D. and Tatsumoto M. (1975a) U-Th-Pb systematics of selected samples from Apollo 17, Boulder I, Station 2. *The Moon* 14, 463-471.
- Palme H., Baddenhausen H., Blum K., Cendales M., Dreibus G., Hofmeister H., Kruse H., Palme C., Spettel B., Vilcsik E. and Wanke H. (1978) New data on lunar samples and achondrites and a comparison of the least fractionated samples from the earth, the moon, and the eucrite parent body. *Proc. 9th Lunar Planet. Sci. Conf.* 25-57.
- Reed V.S. and Wolfe E.W. (1975) Origin of the Taurus-Littrow Massifs. *Proc. 6th Lunar Sci. Conf.* 2443-2462.
- Ryder G. (1983) Nickel in olivines and parent magmas of lunar pristine rocks. In *Workshop on Pristine Highlands Rocks and the Early History of the Moon* (Longhi and Ryder, eds.) LPI Tech Rept. 83-02. The Lunar and Planetary Institute, Houston, 66-68.
- Ryder G. (1984) Most olivine in the lunar highlands is of shallow origin (abs). *LPS XV*, 707-708.
- Ryder G. (1992) A distinct poikilitic impact melt rock from the Apollo 17 landing site that is not from the Serenitatis melt sheet. (abs) *Meteoritics* 27, 284.
- Ryder G. (1993a) The Apollo 17 samples: The massifs and landslide. In *Workshop on Geology of the Apollo 17 Landing Site*. LPI Tech. Rpt. 92-09, 48-49.
- Ryder G. (1993b) Impact melt breccias at the Apollo 17 landing site. In *Workshop on Geology of the Apollo 17 Landing Site*. LPI Tech. Rpt. 92-09, 49-50.
- Ryder G. (1993c) **Catalog of Apollo 17 rocks: Stations 2 and 3.** Curators Office JSC#26088.
- Ryder G., Stoeser D.B., Marvin U.B. and Bower J.F. (1975a) Lunar granites with unique ternary feldspars. *Proc. 6th Lunar Sci. Conf.* 435-449.
- Ryder G., Stoeser D.B., Marvin U.B., Bower J.F. and Wood J.A. (1975b) Boulder 1, Station 2, Apollo 17: Petrology and petrogenesis. *The Moon* 14, 327-357.
- Ryder G. and Spudis P. (1980) Volcanic rocks in the lunar highlands. *Proc. Conf. Lunar Highlands Crust*, 353-375. *Geochim. Cosmochim. Acta suppl. 12*, Lunar Planetary Inst.
- Ryder G., Norman M.D. and Score R.A. (1980a) The distinction of pristine from meteorite-contaminated highlands rocks using metal compositions. *Proc. 11th Lunar Planet. Sci. Conf.* 471-479.
- Ryder G., Norman M.D. and Score R.A. (1980b) Ni, Co content of metal grains for the identification of indigenous rocks (abs). *LPS XI*, 968-970.
- Ryder G., Norman M.D. and Taylor G.J. (1997) The complex stratigraphy of the highland crust in the Serenitatis region of the Moon inferred from mineral fragment chemistry. *Geochim. Cosmochim. Acta* 61, 1083-1105.
- Schaeffer O.A., Warasila R. and Labotka T.C. (1982a) Ages of Serenitatis breccias: Lunar breccias and soils and their meteoritic analogs. *LPI Tech. Rept.* 82-02, 123-125.
- Schaeffer O.A., Warasila R. and Labotka T.C. (1982b) Ages of Serenitatis breccias (abs). *LPS XIII*, 685-686.
- Schmitt H.H. (1975) Geological model for Boulder 1 at Station 2, South Massif, Valley of Taurus-Littrow. *The Moon* 14, 491-504.
- Simonds C.H., Phinney W.C. and Warner J.L. (1974) Petrography and classification of Apollo 17 non-mare rocks with emphasis on samples from the Station 6 boulder. *Proc. 5th Lunar Sci. Conf.* 337-353.
- Stoeser D.B., Wolfe R.W., Marvin U.B., Wood J.A. and Bower J.F. (1974b) Petrographic studies of a boulder from the South Massif (abs). *LS V*, 743-745.
- Stoeser D.B., Wolfe R.W., Wood J.A. and Bower J.F. (1974c) Petrology and petrogenesis of boulder 1. In *Interdisciplinary Studies of Samples from Boulder 1, Station 2, Apollo 17. Volume 1*, Consortium Indomitable. Smithsonian Astrophysical Observatory. *Also Lunar Science Institute Cont. no. 210D*, 35-109.
- Stoeser D.B., Marvin U.B. and Bower J.F. (1974d) Petrology and petrogenesis of boulder 1. In *Interdisciplinary Studies of Samples from Boulder 1, Station 2, Apollo 17. Volume 2*, Consortium Indomitable. Smithsonian

Astrophysical Observatory. *Also* Lunar Science Institute
Cont. no. 21 ID, 1-59.
72215 72235 72275

Stoeser D.B., Ryder G. and Marvin U.B. (1975) Lunar granite clasts with unique ternary feldspars (abs). LS VI, 780-782.

Spudis P.D. and Ryder G. (1981) Apollo 17 impact melts and their relation to the Serenitatis basin. *In* Proc. of the Conf. on Multi-Ring Basins. Proc. Lunar Planet. Sci. 12A - Geochim. Cosmochim. Acta, Suppl. 15. Pergamon Press. 133-148.

Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and volatile elements. Proc. 10th Lunar Planet. Sci. Conf. 2107-2130.

Wolfe E.W. et al. (1981) The geologic investigation of the Taurus-Littrow valley: Apollo 17 landing site. US Geol. Survey Prof. Paper 1080.